

GROUND WATER LEVEL BULLETIN

NOVEMBER- 2024

HIMACHAL PRADESH



ABSTRACT

Ground water level Scenario during Nov-2024 highlighting the findings, status of ground water level in different aquifers and its seasonal, annual and decadal comparison.

INTRODUCTION

Groundwater bulletin is prepared by CGWB depicting changes in groundwater regime of the country through different seasons. It is an effort to obtain information on groundwater levels through representative monitoring wells. The important attributes of groundwater regime monitoring are groundwater level.

The natural conditions affecting the groundwater regime involve climatic parameters like rainfall, evapotranspiration etc., whereas anthropogenic influences include pumpage from the aquifer, recharge due to irrigation systems and other practices like waste disposal etc.

Groundwater levels are being measured by Central Ground Water Board four times a year during January, May, August and November. The regime monitoring started in the year 1969 by Central Groundwater Board. A network of 25437 observation wells called **National Hydrograph Network Stations (NHNS)**, as on 30.04.2023, located all over the country is being monitored.

STUDY AREA

Himachal Pradesh is located between the north latitude $30^{\circ}22'40''$ & $33^{\circ}12'40''$ and east longitude $75^{\circ}47'55''$ & $79^{\circ}04'20''$. It falls in Survey of India topographic sheets Nos. 52D, 52H, 52L, 53A, 53B, 53F, 53E and 53I and covers an area of 55,673 sq km. It is one of the predominantly hilly states in India, which lies in the western Himalayas. The length of Himachal Pradesh is about 355 km i.e., from northwestern part of Chamba to southeastern part of Kinnaur. The breadth of the state is about 270 km i.e., from western part of Una to northeast part of Lahaul and Spiti District. The state is bounded by the state of Jammu & Kashmir in north, Punjab state in west, Haryana state in south and Uttarakhand state in southeast and shares an international border with China (Tibet) in northeast.

Administratively, the state is divided in 12 Districts, 76 Tehsils, 34 Sub-Tehsils and 78 Blocks. There are 20,690 villages, 3,226 Gram Panchayats, 59 towns, 28 Nagar Panchayats and 25 Nagar Parishads including Municipal Corporations. Lahaul & Spiti is the largest and Hamirpur is the smallest District of the state with geographical area of 12,835 and 1,118 sq km respectively.

The state has a population of 68,64,602 persons (Census 2011) having an average population density of 128 person per sq km. The male population in the state is 34,81,873 persons and female population is 33,82,729. The rural

and urban population in the state is 61,76,050 and 6,00,552 persons respectively. The density of population in the state varies from as low as 2 persons/sq.km in Lahaul and Spiti District to 406 persons/sq km in Hamirpur District as compared to the state average of 128 persons/sq km.

Himachal region presents an intricate mosaic of mountain ranges, hills and valleys with altitude ranging from 350 m to 6500 m amsl. The Dhauladhar range looks in supreme majesty over the Kangra valley while the Pir Panjal, the Great Himalaya and the Zaskar ranges guard over Chamba, Lahaul & Spiti, Kullu and Kinnaur Districts. The mountain slopes are covered with forests and meadows. The valleys below are interspersed with numerous streams, fields and quiet homesteads. There is general increase in elevation from east to west and from south to north. The physiographic divisions from south to north are the Outer Himalayas also known as Siwaliks (350 to 1500 m amsl), the Lesser Himalayan Range (1500-5000 m amsl), Great Himalayan Range (5000 – 6000 m amsl) and Zaskar Range (> 6000 m amsl)

Himachal state has a unique distinction of having drainage systems of both the Indus and the Ganga basin. The major river systems of the region are the Chandra-Bhaga or the Chenab, the Ravi, the Beas, the Satluj and the Yamuna. The catchments of the rivers are fed by snow and rainfall, and are protected by fairly extensive cover of natural vegetation. Major rivers of the Indus River basin are the Chenab, the Ravi, the Beas and the Satluj. The Yamuna is the only river contributing water to Ganga basin.

The southwestern monsoon contributes about 70% of rain fall during monsoon period from July to September and about 30% occurs during non-

monsoon period due to western disturbances and in the form of thunder storm. Generally, rainfall increases from south to north. Western disturbances also shower rainfall in winters. Beyond Kullu towards Lahaul & Spiti and Kinnaur, rain fall decreases due to rain shadow effect. Spiti is the driest area with 50 mm rainfall because of being enclosed by High Mountain from all sides. The average annual rainfall in the Districts of the state varies from about 600 mm in Lahaul & Spiti to more than 2400 mm in Kangra.

GROUND WATER LEVEL MONITORING

Central Ground Water Board, Northern Himalayan Region, is monitoring changes in groundwater regime in Himachal Pradesh state on quarterly basis continuously. This is facilitated by a network of monitoring stations in the State located in diverse hydrogeological and geomorphic units. The number of operational wells till Nov 2024 was 223 which include 137 dug wells, 59 piezometers and 28 springs. In Nov 2024, 224 wells which include 137 dug wells, 60 piezometers and 27 springs. The district-wise breakup of the water level monitoring stations is given in **Table-1**.

Sl. No	Name of the District	Number of GW Monitoring Stations (NOV-2024)				
		Dug Well	Piezometer	Hand pump	Spring	Total
1	Hamirpur	6	3	-	-	9
2	Bilaspur	4	8	-	-	12
3	Kangra	46	19	-	2	67
4	Kullu	2	3	-	-	5
5	Mandi	10	-	-	4	14
6	Sirmaur	19	12	-	-	31
7	Solan	14	5	-	-	19
8	Una	34	8	-	-	42
9	Chamba	2	2	-	21	25
	TOTAL	137	60	-	27	224

Table-1 District-wise breakup of the water level monitoring stations

RAINFALL

The rainfall data collected and compiled from weekly and monthly weather reports from India Meteorological Department were used to analyze the rainfall for the period June 2024 – September 2024. Table-2 gives the district-wise rainfall data for the period June 2024 – September 2024.

Month- June 2024				
S.No	District	Act (mm)	Nor (mm)	Dep (%)
1	Bilaspur	44.4	102.1	-57
2	Chamba	40.1	121.8	-67
3	Hamirpur	39.6	114.1	-65
4	Kangra	60.7	181.4	-67
5	Kinnaur	16.1	41.8	-61
6	Kullu	43.4	86.9	-50
7	Lahaul & Spiti	25	47.8	-48
8	Mandi	73.8	171.5	-57
9	Shimla	80.3	110.4	-27
10	Sirmaur	76.5	165.8	-54
11	Solan	89.4	137.3	-35
12	Una	54.9	103.8	-47
13	Sub- Division HP	46.2	101.1	-54

Month- July 2024				
S.No	District	Act (mm)	Nor (mm)	Dep (%)
1	Bilaspur	204.5	272.2	-25
2	Chamba	196.2	305.7	-36
3	Hamirpur	207.9	328.5	-37
4	Kangra	581.5	589.3	-1
5	Kinnaur	38.3	65.9	-42
6	Kullu	161.4	184	-12
7	Lahaul & Spiti	12.1	131.5	-91
8	Mandi	352.6	386.5	-9
9	Shimla	195.3	210.2	-7
10	Sirmaur	254	437	-42

11	Solan	142.4	303.3	-53
12	Una	176.8	329	-46
13	Sub- Division HP	180.5	255.9	-29

Month- Novust 2024				
S.No	District	Act (mm)	Nor (mm)	Dep (%)
1	Bilaspur	404.8	316.8	28
2	Chamba	239.3	291.7	-18
3	Hamirpur	320.5	400.6	-20
4	Kangra	654.8	631.5	4
5	Kinnaur	52.6	77.6	-32
6	Kullu	144.5	180.2	-20
7	Lahaul & Spiti	43.4	117.6	-63
8	Mandi	421.6	395.3	7
9	Shimla	300	196.4	53
10	Sirmaur	522.9	402.1	30
11	Solan	271.6	287.9	-6
12	Una	305.3	372.2	-18
13	Sub- Division HP	243.6	256.8	-5

Month- September 2024				
S.No	District	Act (mm)	Nor (mm)	Dep (%)
1	Bilaspur	182.8	128	43
2	Chamba	91.9	134.2	-31
3	Hamirpur	128.4	130	-1
4	Kangra	274.2	220.2	25
5	Kinnaur	87.2	62.5	40
6	Kullu	102.3	96.9	6
7	Lahaul & Spiti	23.1	86	-73
8	Mandi	220.1	144.2	53
9	Shimla	145.9	110.5	32
10	Sirmaur	312.1	178.9	74
11	Solan	172.9	145.8	19
12	Una	111	148	-25
13	Sub- Division HP	125.3	120.6	4

Table-2 District-wise Rainfall from 01.06.2024 to 30.09.2024

LEGEND: ■ L. EXCESS (+60% OR MORE) ■ EXCESS (+20% TO +59%) ■ NORMAL (+19% TO -19%)
■ DEFICIENT (-20% TO -59%) ■ L. DEFICIENT (-60% TO -99%) ■ NO RAIN (-100%) NO DATA

INDIA METEOROLOGICAL DEPARTMENT
MC SHIMLA

Rainfall (in mm)/ Departures (in %) from the Long Period Averages
for Districts in HIMACHAL PRADESH

PERIOD: 01.06.2024 - 30.09.2024



LEGEND: ■ L. EXCESS (+60% OR MORE) ■ EXCESS (+20% TO +59%) ■ NORMAL (+19% TO -19%)
■ DEFICIENT (-20% TO -59%) ■ L. DEFICIENT (-60% TO -99%) ■ NO RAIN [-100%] ■ NO DATA

A perusal of the DTWL map for November 2024 shows that the shallow water level areas of less than 2 m observed in southern part of Kangra Palampur valley and southern part of Balh valley. Water level of 2-5 m & 5-10 m bgl is observed in major part of Kangra Palampur valley, Indaura-Nurpur valley, Balh valley, Una Valley, Nalagarh valley Paonta valley respectively. 10-20 m bgl water levels shown in Una, Nalagah, Kangra-Palampur valley and Paonta valley only. Deeper water level more than 20 m is confined mainly in northern part of Paonta valley in Sirmaur District, northern part of Nalagarh valley of Solan District and northern part of Una valley.

Season	No of well analysed	Number and Percentage of wells showing depth to water level (mbgl) in the range of											
		0 to 2		2 to 5		5 to 10		10 to 20		20 to 40		> 40	
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Nov-24	197	41	20.81	56	28.43	49	24.87	26	13.20	19	9.64	6	3.05

Depth to Water Level Distribution of Percentage of Observation Wells Pre -monsoon 2024

DEPTH TO WATER LEVEL (NOV-2024)

The depth to water level, recorded during Nov 2024 ranged between 0.08 m (Bilaspur District) and 95.32 m bgl (Tikkri (Pz) Solan District). Out of 197 stations monitored, the majority of 172 NHS (87.31%) recorded DTWL, in the range between 2 - 20 m bgl. 41 stations (20.81%), recorded shallow water levels, less than 2 m bgl and 25 stations (12.69%), recorded deep water levels, more than 20 m bgl in the state.





Fig:1 Depth to Water Level Nov-2024


SEASONAL WATER LEVEL FLUCTUATION

To study the effect of monsoon and subsequent utilization for various needs like agriculture, irrigation and domestic etc, changes in water level are studied and are discussed below.

May 2024 to Nov 2024

Seasonal fluctuation of water level was analyzed for 197 stations for the period May 2024 – Nov 2024. Out of the 197 stations, 148 stations have shown rise in water level and remaining 49 stations have shown fall in water level.

The minimum rise in water level of 0.02 m was observed in Una and Hamirpur District and the maximum rise 12.87 m was noticed in Kangra District (Sapri Pz). Out of the 148 stations which have shown rise in water level, 122 stations show rise between the range of 0 to 2 m, 12 stations between 2 to 4 m and remaining 14 stations show rise more than 4 m.



Comparison	Season	No of wells analysed	Number and percentage of wells in different depth range											
			Rise						Fall					
			0 to 2	%	2 to 4	%	> 4	%	0 to 2	%	2 to 4	%	> 4	%
Seasonal	May-24-Nov-24	197	122	61.93	12	6.09	14	7.11	43	21.83	4	2.03	2	1.02

ANNUAL WATER LEVEL FLUCTUATION

Annual fluctuation in water level of GWMS during different monitoring period was analysed and discussed below.

November 2023 to November 2024

Annual fluctuation of water level, has been worked out by comparing depth to water level of Nov 2023, with Nov 2024 and the data presented its frequency distribution in various rise and fall ranges.

Out of the 197 stations analysed, 82 stations (41.63%) have shown rise in water level ranging from 0.02 (Mandi District) to 4.65 m (Kangra District).

Out of 82 stations which have shown rise in water level, 78 stations (95.12%) show rise between the range of 0 to 2 m, 2 station (2.44%) has shown rise between 2 to 4 m and 2 station (2.44%) shown rise more than 4 m.

Similarly, for 115 stations which have shown fall in water level, 87 stations (75.65%) show fall between the range of 0 to 2 m, 21 stations (18.26%) have shown fall between 2 to 4 m and 7 stations (6.09%) has shown fall more than 4 m.

A perusal of map of annual fluctuation of November 2023 to November 2024 showing fall in water levels in Nurpur Indora valley and central part of Una valley, Paonta Valley, Balh Valley and Nalagarh Vally. Similarly rise in water level 0-2 m is noticed along the fringe areas of all monitoring valleys except Kullu Valley and Balh Valley (Mandi District).




Fig:2 Seasonal fluctuations for the period May 2024 – Nov 2024

Comparison	Season	No of wells analysed	Number and percentage of wells in different depth range											
			Rise						Fall					
			0 to 2	%	2 to 4	%	> 4	%	0 to 2	%	2 to 4	%	> 4	%
Annual	Nov-24	197	78	39.59	2	1.02	2	1.02	87	44.16	21	10.66	7	3.55





Fig:3 Annual fluctuations for the period Nov 2023 – Nov 2024

DECADAL FLUCTUATIONS

The decadal variations were analyzed considering the decadal average of water level and the water level for the respective period.


Decadal average of Nov (2014-2023) to Nov 2024


Decadal water level fluctuation has been worked out by comparing water level data of Nov 2024 with the average mean of 10 years' water level data of Nov (2014-2023) and is presented frequency distribution in various ranges.

A perusal shows that out of 104 stations analyzed, 37 stations (35.58%) have shown rise and 67 stations (64.42%), have shown fall in water level. Out of 37 stations 29 stations (78.38%) are showing rise in water level between 0 to 2 m, 5 stations (13.51%) between 2 to 4 m and 3 stations (8.11%), more than 4 m. Out of 67 stations, 59 stations (88.06%) show fall in water level between 0 to 2 m, 4 stations (5.98 %) between 2 to 4 m and 4 stations (5.98%) more than 4 m. A minimum rise in water level of 0.012 m was noticed in Mandi Districts and the maximum rise of 7.11 m is noticed in Kangra District. Similarly, the minimum fall of 0.01 m is noticed in Kangra District & maximum fall of 9.01 m is noticed in Kangra District.

A perusal of map of Decadal average of November (2014-2023) to November 2024 reveals rise in water level less than 2 m is shown in whole part of Kangra- Palampur valley & Indaura valley of Kangra District except a few places, major part of Nalagarh valley, Balh valley, a couple of places in Paonta valley. The fall between 2 to 4 m was noticed in, Una valley and Paonta valley. Similarly, rise is noticed in all the valleys from 0-2 m .

Comparison	Season	No of wells analysed	Number and percentage of wells in different depth range											
			Rise						Fall					
			0 to 2	%	2 to 4	%	> 4	%	0 to 2	%	2 to 4	%	> 4	%
Decadal	Nov-24	104	29	27.88	5	4.81	3	2.88	59	56.73	4	3.85	4	3.85





SUMMARY

As a component of the National Ground Water Monitoring Programme, the CGWB, NHR, Dharamshala conducts monitoring of the ground water conditions on a quarterly basis: in January, Pre-monsoon (May), August and Post-monsoon (November). Additionally, a yearly assessment of ground water quality is performed in May and Nov-2024, the Northern Himalayan Region of the Central Ground Water Board monitored 137 dug wells, 60 piezometers and 27 springs. This comprehensive effort aims to portray the variations in the state's ground water conditions across different aquifers.

In Nov 2024, 87.31% of the state's area exhibited a depth to water level within 20 meters below ground level. Deeper water levels of more than 20 m cover 12.69 % area of the State covering mainly Una, Sirmaur, Solan, Kangra, districts.

Annual water level comparison with previous year Nov-2023 to Nov-2024 has shown that about 41.63% area of the state experienced rise in annual water level fluctuation. 35.57% of the area experienced rise in decadal mean water level of 2014-2023 with respect to Nov, 2024.

Fig. 4: Decadal water level fluctuation, Decadal mean Nov (2014-2023) Vs Nov-2024