




### Percentage of Wells In Different Water Level Ranges In Unconfined Aquifers (Aug, 2024)



- 0-2 mbgl ■ 2-5 mbgl ■ 5-10 mbgl
- 10-20 mbgl ■ 20-40 mbgl ■ >40 mbgl



## ABSTRACT

Ground water level Scenario during August-2024 highlighting the findings, status of ground water level in different aquifers and its seasonal, annual and decadal comparison.

**CGWB, NORTH WESTERN REGION, CHANDIGARH**

# GROUND WATER LEVEL BULLETIN

PUNJAB STATE AND CHANDIGARH UT

## 1.0 INTRODUCTION


Groundwater bulletin is prepared by CGWB depicting changes in groundwater regime of the country through different seasons. It is an effort to obtain information on groundwater levels through representative monitoring wells. The important attributes of groundwater regime monitoring are groundwater level.

The natural conditions affecting the groundwater regime involve climatic parameters like rainfall, evapotranspiration etc., whereas anthropogenic influences include pumpage from the aquifer, recharge due to irrigation systems and other practices like waste disposal etc.

Groundwater levels are being measured by Central Ground Water Board four times a year during January, June, August and November. The regime monitoring started in the year 1969 by Central Groundwater Board. A network of 1874 observation wells called **National Hydrograph Network Stations (NHNS)**, as on 30.04.2024, located all over the country is being monitored.

## 2.0 STUDY AREA

The Punjab State is located between North latitudes  $29^{\circ} 32'$  and  $32^{\circ} 28'$  and East longitudes  $73^{\circ} 50'$  and  $77^{\circ} 00'$ . The total geographical area of the state is 50476 sq. km. It is surrounded by the states of Himachal Pradesh in the northeast, Jammu and Kashmir in the north and Haryana and Rajasthan in the south and southwest respectively. The State has a flat alluvial plain except a narrow belt along the southwestern parts where stable sand dunes are seen dotting the landscape. The area occupied by the mountains (Himalayan foothills) in northeast, is about 1243 sq km. Perennial Rivers Sutlej, Beas, Ravi and ephemeral river Ghaggar drain the state. It has a vast network of canal system. With the inception of canals, the fertile land of the State started converting into green fields and experienced spectacular achievement in agricultural production with emphasis on cultivation of paddy and high yielding varieties of crops, as a consequence demand for water increased manifold resulting in over exploitation of ground water resources.



**Figure-1: Map showing major aquifers and Hydrogeology of Punjab State & Chandigarh UT**

The State has been divided into four main divisions viz. Jalandhar, Patiala, Ferozpur and Faridkot, which are further sub-divided into 23 districts, which are further divided into 77 sub-divisions /tehsils and 146 community development.

The State forms a part of vast Indo-Genetic alluvial plain. Physiographically, the State can be divided into seven distinct units, which run parallel to each other.

**I Hilly area:** Siwalik Hills on the north and northeastern part.

**II Eroded hills with flat land (Plateau):** forms top of hills.

**III Intermountain valleys**

**IV Piedmont area:** (Kandi zone) immediately southwest of hills.

### 3.0 GROUND WATER LEVEL MONITORING

The Central Ground Water Board, North Western Region, has established 521 Ground water observation wells in Punjab State and Union Territory of Chandigarh for monitoring water level. As on 31.3.2024 there were 498 Ground Water Observation Wells of CGWB in Punjab which include 114 dug wells and 384 piezometers for monitoring shallow & deeper aquifers in Punjab. There are 23 observation wells in Chandigarh, 1 dug wells and 22 Piezometers for monitoring shallow & deeper aquifers.

About 80% of the Ground water observation wells fall in the canal command areas of various canal systems, the areas falling out of the major command is part of Pathankot, Hoshiarpur, Nawanshahr, Ropar and SAS Nagar districts, parts of Gurdaspur, Jalandhar and Ludhiana districts. The district wise details of Ground water observation wells are given in Table 1 and location of these Ground water observation wells is shown in Figure 2.

**V. Sirowal Zone:** lies further southwest of Kandi area which merges with the alluvium of Ravi, Beas, Sutlej and Ghaggar rivers. 2

**VI. Alluvial plains:**

- Active/recent flood plains include meanders and present flood plains.
- Abandoned flood plains include terraces of rivers, abandoned during Recent age.
- Bar upland areas: Higher elevated land which remained beyond the reach of rivers but are composed of ancient river channels deposits (older alluvium) plains.

**VII. Sand Dunes:** Covering southwest part




Figure- 2: Map showing locations of monitoring wells (NHNS) in Punjab

S. No.	District	Dug Well	Tube Well	Total
1	Amritsar	3	14	17
2	Barnala	0	7	7
3	Bathinda	14	20	34
4	Faridkot	11	10	21
5	Fatehgarh Sahib	3	19	22
6	Fazilka	7	18	25
7	Firozpur	2	19	21
8	Gurdaspur	10	21	31
9	Hoshiarpur	13	17	30
10	Jalandhar	2	43	45
11	Kapurthala	0	18	18
12	Ludhiana	3	31	34
13	Mansa	4	15	19
14	Moga	0	14	14
15	Muktsar	7	7	14
16	Pathankot	13	0	13
17	Patiala	4	32	36
18	Rupnagar	7	8	15
19	Sangrur	1	32	33
20	Sas Nagar	8	11	19
21	Sbs Nagar	0	9	9
22	Tarn Taran	2	19	21
23	Chandigarh UT	1	22	23
<b>Grand Total</b>		<b>115</b>	<b>406</b>	<b>521</b>

**Table 1: District-wise distribution of water level monitoring stations**

#### 4.0 GROUND WATER LEVEL SCENARIO (AUGUST, 2024)

##### 4.1 SHALLOW AQUIFER (UNCONFINED)

###### 4.1.1 DEPTH TO WATER LEVEL

###### Depth To Water Level in Unconfined Aquifer (August, 2024)

The behavioral pattern of water level in August 2024 along with depth to water level map (Fig.3) is discussed below.

The depth to water level lies between 0.27 mbgl in Faridkot district and 60.36 mbgl in Hoshiarpur district. Very shallow water levels of 0-2 m (causing water logging) occur in 6.88% of wells and 1.36% area of the state in isolated patches in Muktsar, Faridkot and Fazilka districts. Shallow water levels of 2-5 m have been observed in 17% of the wells and 10.32% of the total area that lies in south western parts of state i.e Muktsar, Fazilka, Faridkot and parts of Gurdaspur, Pathankot, Rupnagar and Ludhiana districts. These are mainly canal command areas and use canal water for their agricultural needs. The water levels between 5-10 m are observed in Fazilka, Faridkot, Muktsar, Ferozepur, Bathinda, Mansa, Gurdaspur, Pathankot, Hoshiarpur, SBS Nagar, Patiala and Rupnagar districts. About 17% of wells and 14.15% of the area fall in this range. Moderately Deep water levels (10-20 m) are observed in 21.86% wells covering about 25.37% area of the State Pathankot, Gurdaspur, Amritsar, SBS Nagar, Hoshiarpur, Jalandhar, Tarn Taran, Kapurthala, Moga, Ludhiana, Fazilka, Ferozpur, Faridkot, Bathinda, Mansa, Rupnagar & SAS Nagar districts. Deep water levels (20-40 m) are observed in parts of Amritsar, Taran taran, Jalandhar, Kapurthala, SBS Nagar, Hoshiarpur, Bathinda, Mansa, Moga, Ludhiana, Fatehgarh Sahib, Patiala, Sangrur, Barnala and SAS Nagar districts and observed in 27.94% wells covering about 37.59% area of the state. Very deep water levels (>40 m) are observed in 9.31% wells as patches in Sangrur, Barnala, Malerkotla, Patiala & SAS Nagar districts covering 11.20% area of the State.




Figure-3: Depth to Water Level Map Unconfined Aquifer, August 2024

###### Percentage of Wells In Different Water Level Ranges In Unconfined Aquifers (Aug, 2024)






Figure-4: Percentage of wells in different water level ranges in unconfined aquifer.



**Wells Showing Rise & Fall In Seasonal Fluctuation  
(Jun, 2023 to Aug, 2024)**



**Figure-5: Percentage of wells showing rise and fall in WL in unconfined aquifer(June 2024 to August 2024)**



**Figure-6: Seasonal water level fluctuation in unconfined Aquifer (June 2024 to August 2024)**

#### 4.1.2 SEASONAL FLUCTUATION IN WATER LEVEL

##### Seasonal Fluctuation of Water Level in Unconfined Aquifer (June 2024 to August 2024)


Water level data of August 2024 when compared with previous measurement data i.e. January 2024 is termed as seasonal water level fluctuations. The behavioral pattern of this seasonal fluctuation is discussed below. The map depicting seasonal water level fluctuations is shown in Fig.6.

##### Rise in Water Levels:


The water level rise has been recorded in 39.53% of wells monitored and covering 72.83% area of the State. Water level rise in the range of 0-2 m is observed in 30.48% of wells and 51.27% of the area. Water level rise 2-4m is observed in 6.67% of the wells & 19.85% area. Water level rise of >4m is observed in 2.38% wells and in 1.71% area as isolated patches in Nawanshahar, Kapurthala, Ferozpur and Patiala districts.

##### Fall in Water Levels:

The seasonal fluctuation shows that there is a general decline of water levels in 60.47% of wells monitored and covering 27.17% area of the State. The decline has been observed in all districts except some isolated patches scattered over the state. Water level decline in the range of 0-2 m is observed in 38.57% of wells and 23.61% of area. Water level decline in the range of 2-4 m is observed in 17.14% of wells and 2.35% of area. Water level decline of >4m is observed in 4.76% of wells and less than 1.21% of area as isolated patches in Kapurthala, Bathinda, Muktsar, Pathankot & Faridkot districts.



**Figure-7: Percentage of wells showing rise and fall in WL in unconfined aquifer  
(August 2023 to August 2024)**



**Figure-8: Annual water level fluctuation in unconfined aquifer(August 2023 to  
August 2024)**

**4.1.3 ANNUAL FLUCTUATION IN WATER LEVEL**

**Annual Fluctuation of Water Level in Unconfined Aquifer (August 2023 to August 2024)**



In order to know the impact of rainfall and ground water withdrawal during last one year, annual water level fluctuations for period August 2023 and August 2024 are calculated. The behaviour of annual fluctuations is discussed in the following paragraph and depicted in Fig.8.

**Rise in Water Levels:**

The water level rise has been recorded in 24% of wells monitored and covering 25.86% area of the State. Water level rise in the range of 0-2 m is observed in 21.5% wells and less than 1% of area. Water level rise 2-4m is observed in 1.5% wells and 1.94% of area. The water level rise of >4m is observed in 1% wells and less than 1% of area as isolated patches in Tarn Taran, Faridkot, Hoshiarpur and Chandigarh UT.

**Fall in Water Levels:**

The annual fluctuation depicts general decline of water levels in 76% of wells monitored and covering 74.14% area of the State. The decline has been observed in all districts of the state. Water level decline the range of 0-2 m is observed in 64.5% of wells and 66.58% of the area. Water level decline in the range of 2-4 m is observed in 9.5% of wells and 5.52% of the area. Whereas, the water level decline of >4m is observed in 2% of wells and 2.04% of the area during the period, as isolated patches in Faridkot, Moga, Sangrur, Barnala and Kapurthala districts.



#### 4.1.4 PAST TWO YEAR FLUCTUATION IN WATER LEVEL

### Past Two Year Fluctuation of Water Level in Unconfined Aquifer (August 2022 to August 2024)

In order to know the impact of rainfall and ground water withdrawal during last two years, past two year water level fluctuations for period August 2022 and August 2024 are calculated. The behaviour of annual fluctuations is discussed in the following paragraph and depicted in Fig.10.


### Rise in Water Levels

The water level rise has been recorded in 34.4% of wells monitored and covering 9.93% area of the State. Water level rise in the range of 0-2 m is observed in 26.34% wells and 8.53% of area. Water level rise 2-4m is observed in 6.45% wells and 1.24% of area. The water level rise of >4m is observed in 1.61% wells and less than 1% of area as isolated patches in Patiala, Sangrur, Barnala and Moga districts.


## Fall in Water Levels

The annual fluctuation depicts general decline of water levels in 65.6% of wells monitored and covering 90.07% area of the State. The decline has been observed in all districts of the state. Water level decline the range of 0-2 m is observed in 50% of wells and 19.91% of the area. Water level decline in the range of 2-4 m is observed in 12.37% of wells and 69.31% of the area. Whereas, the water level decline of >4m is observed in 3.23% of wells and less than 1% of the area during the period, as isolated patches in Patiala and Sangrur districts.





**Figure-11: Percentage of wells showing rise and fall in WL in unconfined Aquifer (Decadal Mean August (2014-2023) to August 2024)**



**Figure-12: Water level fluctuation in unconfined Aquifer(Decadal Mean August (2014-2023) to August 2024)**

#### 4.1.5 DECADAL FLUCTUATION IN WATER LEVEL

##### Decadal Fluctuation of Water Level in Unconfined Aquifer (Decadal Mean August (2014-2023) to August 2024)

Changes in water level behaviour since last one decade are determined using decadal mean data. Water level mean of past one decade (2014-2023) for each ground water observation well is computed and compared with the respective water level data of August 2024. The behaviour of water level over the period under reference is discussed in paragraph below along with Fig.12.


##### Rise in Water Levels:

The decadal mean fluctuations show that rise in 24.57% of observation wells monitored covering about 16.74% area of the state. Water level rise in the range of 0-2 m is observed in 24.69% of wells and 15.37% of the area. Water level rise of 2-4m is observed in 1.65% of wells and 1.12% of the area. Water level rise of >4m is observed in 1.23% of wells and less than 1% of the state area as isolated patches in Hoshiarpur district.

##### Fall in Water Levels:

The decadal mean fluctuations show that decline in 72.43% of observation wells monitored covering about 82.26% area of the state. The decline has been observed in all districts of the state. The decline of 0-2 m has been observed in about 40.33% of wells and 45.41% of area. Water level decline of 2-4 m is observed in 16.46% of the wells and 20.31% of the area. Water level decline of >4m is observed in 15.64% of the wells and 17.54% of area during the period, in Barnala, Sangrur, Malerkotla, Moga, SBS Nagar, Jalandhar, Kapurthala, Patiala and Bathinda districts.

### Percentage of Wells In Different Water Level Ranges In Confined Aquifers (Aug, 2024)




**Figure-13: Percentage of wells in different water level ranges in Confined/Semi-Confined aquifer.**


## 4.2 DEEPER AQUIFER (CONFINED/ SEMI-CONFINED)

### 4.2.1 DEPTH TO PIEZOMETRIC LEVEL

The behavioral pattern of water level in August 2024 along with depth to water level map (Fig. 14) is discussed here. The depth to water level lies between 0.98mbgl in Gurdaspur district and 67.16mbgl in Hoshiarpur district. Very shallow water levels of 0-2 m (causing water logging) does occur in 5.06% of the wells and 1.86% of the state's area. Shallow water levels of 2-5 m have been observed in 2.53% of the wells and 17.04% of the total area. The water levels between 5-10 m are about 2.53% of wells and 11.94% of the area fall in this range. Moderately Deep-water levels (10-20 m) are observed in 18.99% wells covering about 12.72% area of the State. Deep water levels (20-40 m) are observed in 44.3% wells covering about 39% area of the state. Very deep-water levels (>40 m) are observed in 26.58% of the wells covering 17.44% area of the State.



**Figure-14: Depth to Water Level Map Confined/Semi-Confined Aquifer, August 2024**



**Figure-15: Annual water level fluctuation in Confined/Semi-Confined aquifer (August 2023 to August 2024)**

#### 4.2.2 ANNUAL FLUCTUATION IN WATER LEVEL

##### Annual Fluctuation of Water Level in Confined Aquifer (August 2023 to August 2024)

In order to know the impact of rainfall and ground water withdrawal during last one-year, annual water level fluctuations for period August 2023 and August 2024 for the confined aquifers and semi-confined aquifers are calculated. The behavior of annual fluctuations is discussed in the following paragraph and depicted in Fig 15.

##### Rise in Water Levels:

The water level rise has been recorded in 32.38% area of the State. Water level rise in the range of 0-2 m is observed in 26.85% of area. Water level rise 2-4m is observed in 4.59% of area. The water level rise of >4m is observed in less than 1% of area as in Mansa, Sangrur, Patiala, Gurdaspur and Hoshiarpur districts.

##### Fall in Water Levels:

The annual fluctuation depicts general decline of water levels in 67.62% area of the State. The decline has been observed in most of the districts of the state. Water level decline the range of 0-2 m is observed in 66.28% of the area. Water level decline in the range of 2-4 m is observed in 1.33% of the area. Whereas, the water level decline of >4m is observed in less than 1% of the area during the period, as isolated patches in Moga, Barnala, Jalandhar and Kapurthala districts.